

stimfruit



0-25-32

GUARANTEED ANALYSIS

| | |
|--|-------|
| Available Phosphate (P ₂ O ₅) | 25 % |
| Soluble Potash (K ₂ O) | 32 % |
| Boron (B) | 0.2 % |
| Manganese (Mn) | 0.1 % |
| 0.1% Chelated Manganese | |
| Molybdenum (Mo) | 0.4 % |
| Zinc (Zn) | 0.1 % |
| 0.1% Chelated Zinc | |

Derived from Dipotassium phosphate, Disodium octaborate tetrahydrate, Manganese EDTA, Zinc EDTA, Sodium molybdate

ACTION

Product developed to increase the size of the fruit due to the effect of potassium and phosphorus

DOSE FOR VEGETABLES

Foliar application: 0.65 oz. per 10 gallon of water
 Root application: 2.0 lb per acre
 Hydroponic: 0.40-0.45 lb per acre.
 Cucurbits: fruits with 7-10 cm diameter. Repeat after 15 days in the greenhouse.
 Pepper, tomato: after the fruit setting. Repeat every 15 days.
 Lettuce, spinach: 15-20 days before harvest
 Strawberry: after the fruit setting. Repeat every 15 days.
 Potato, beet: just before the start of the thickened underground organs and 20 days before harvest.
 Aubergine: consulting technical service.

FOLIAR DOSE FOR APPLES AND PEARS

- First application: 1.3 oz per 10 gallons of water, applied to the 3-4 weeks after petal drop (fruit 8-12 mm in diameter).
- Second application: 0.65 oz per 10 galon of water, 1 month before harvest.

DOSE FOR PEACH, NECTARINE ALMOND AND WALNUT

Apply 1.30 oz per 10 galons of wáter after thinning, can be repeated after 15-20 days before.

FOLIAR DOSE FOR CITRUS TREES

- 1.3 oz per 10 galons of water

Fruit setting phase
 3 applications in 45 days (every 15 days)
 • first: 50 % open petals • second: 80 % fallen petals
 • third: at the end of flowering

Fattening phase
 1-2 applications
 • first: fruit with 2-3 cm • second: 20-25 days after first application

STORAGE

Avoid low temperatures in winter and temperatures above 82°F/28°C. Avoid direct light and long storage periods.

WARNINGS

USED ONLY WHEN REQUIRED.

The recommendations and information given in this label have been carefully studied and tested. However in the utilization of the product could interfere in other factors that are no possible to control, such as mixture preparation, application, climate, etc.). The company guarantees the composition, formulation and content. The user will be responsible of the damages caused by lack of efficacy, toxicity, residues, etc. by total inobservance or partial of the instructions given in the label, keep out of reach of children. Keep away from food, drinks and cattle food.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

CAUTION

Stimfruit is recommended only for use in vegetables (cucurbits, pepper, tomato, aubergine (eggplant), lettuce, spinach, strawberry, potato, and beets), tree fruit (apples, pears, peaches, nectarines, and citrus) and tree nuts (almond and walnuts). The use of Stimfruit on any other crop or at any other timing not listed on the label may result in serious crop injury.

The application of fertilizing materials containing molybdenum (Mo) and Boron (B) may result in forage crops containing levels of molybdenum (Mo) and Boron (B) which are toxic to ruminant animals*

Guaranteed/Distributed by:
Lida Plant Research LLC.
 1444 Fulton Street
 Fresno, CA 93721
 Phone (559) 5003535
 Fax (559) 5003536



Net Weight:
2.2lb/1Kg or 11.02 lb/5 Kg