



signalin[®]

0.5 - 3 - 2

GUARANTEED ANALYSIS

Total Nitrogen (N)	0.5 %
0.5 % Water Soluble Nitrogen	
Available Phosphate (P ₂ O ₅)	3 %
Soluble Potash (K ₂ O)	2 %

Derived from seafood by-product, monopotassium phosphate

ACTION

Signalin is recommended for use as a supplement to improve crop productivity and promote plant vigor in agricultural crops.

STORAGE

Avoid low temperatures in winter and temperatures above 28°C. Avoid direct light and long storage periods.

WARNINGS

USED ONLY WHEN REQUIRED.

The recommendations and information given in this label have been carefully studied and tested. However in the utilization of the product could interfere in other factors that are no possible to control, such as mixture preparation, application, climate, etc.). The company guarantees the composition, formulation and content. The user will be responsible of the damages caused by lack of efficacy, toxicity, residues, etc. by total inobservance or partial of the instructions given in the label. Keep out of reach of children. Keep away from food, drinks and cattle food.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

Manufactured by:



Guaranteed/Distributed by:
Lida Plant Research LLC.
3104 Fredericksburg Way
Modesto, CA 95355
(209) 450-5117

CROPS AND DOSE

Onions, pink and white garlic, leeks, celery, carrots, broccoli and potatoes.

Dose:

- foliar spray: 3 oz. per 10 gal. of water
- drip irrigation: 2 lb/acre

Apply every 15-21 days during the whole crop cycle. The first application should be carried out when the first leaves have appeared so the plants can receive the treatment.

Tomatoes, peppers, courgettes, cucumbers, aubergines, beans, melons (watermelon, cantaloupe, honeydew), strawberries, raspberries, blueberries, blackberries.

Dose:

- foliar spray: 3 oz. per 10 gal. of water

Dose drip irrigation: 2 lb/acre

Apply every 15-21 days during the whole crop cycle. The first application should be carried out in seedbeds or at the time of transplanting.

Net Weight:
2.2lb/1Kg or 11.02 lb/5 kg