



rental

8.5-5-5

GUARANTEED ANALYSIS

Total (N) nitrogen	8.5 %
7.5 % Ureic nitrogen	
1 % Ammoniacal nitrogen	
Available Phosphate (P ₂ O ₅)	5 %
Soluble potash (K ₂ O)	5 %

Derived from Diammonium phosphate, Urea, Dipotassium phosphate.

ACTION

rental helps overcome post-transplant stress states, promotes root development, unlock and improves nutrient uptake and vegetative growth of leaves and fruits.

CAUTION

The use of **rental** on any other crop or at any other timing not listed on the label may result in serious crop injury.

WARNINGS

USE ONLY IN CASE OF NECESSITY.

DO NOT EXCEED THE RECOMMENDED RATE.

The recommendations and information given in this label have been carefully studied and tested. However, when applying the product, there could be interferences with other factors that are not possible to control such as mixture preparation, application, climate, etc. Lida Plant Research LLC. guarantees the composition, formulation and content. The user is responsible for any damages caused by lack of efficacy, toxicity, residues, etc. by partial and/or total un-observance of the instructions given on the label. Keep out of reach of children. Keep away from food, drinks and cattle feed.

STORAGE

Avoid low temperatures in winter and temperatures above 82°F/28°C. Avoid direct sunlight and long storage periods.

Information regarding the contents and levels of metals in this product is available on the internet at: <http://www.aapfco.org/metals.html>

CROPS AND RATES

General concentration for foliar spraying: 0.2 % (2.5 fl. oz per 10 gals of water).

Horticultural crops

· Via irrigation: 3-4 pint /acre ; 3-5 applications throughout the crop cycle, with the first one at the sowing or pre-transplant stage.

Strawberries

· Via irrigation: 3-4 pints/acre ; 2 to 6 applications, 10-15 days after transplant and every 15 days.

· Foliar application: 2 pints/acre ; 4-5 applications every 15 days.

Alfalfa

Foliar application of 2-3 pints/acre after each cut.

Cereal

· Foliar application: 2-3 pints/acre ; 2 applications.

Apply in combination with herbicides to reduce stress and stimulate vegetative growth.

Citrus and fruit trees

· Via irrigation: 4-6 pints/acre; one application at budding stage in spring, at pre-flowering and fruit-setting stages and at budding stage in summer.

Table Grape and Wine Grape

· Via irrigation: 4-6 pints/acre; one application before budding stage, at pre- and post-flowering stages and at the beginning of summer.

Banana trees

Via irrigation: 4-6 pints/acre; 3-4 applications from the formation of the sucker.

Olive trees

· Foliar application: 2-3 pints/acre; One application in spring and another at stone hardening.

Density: 10.6 lbs./gal (1.27 Kg/L) at 68 °F

Net Volume: 2.5 gals (9.46 Liters)

Net Weight: 26.5 lbs. (12.02 Kg)

Batch number: LT20-01457

Production date: July 2020

Expiration date: July 2022



Guaranteed/Distributed by:
Lida Plant Research LLC.
1444 Fulton Street
Fresno, CA 93721
Phone (559) 5003535 . Fax (559) 5003536